

	<b>QMRF identifier (JRC Inventory): Q13-16-0061</b>
	<b>QMRF Title: QSAR for octanol-water partition coefficient for pesticides</b>
	<b>Printing Date: Dec 11, 2019</b>

## 1. QSAR identifier

### 1.1. QSAR identifier (title):

QSAR for octanol-water partition coefficient for pesticides

### 1.2. Other related models:

### 1.3. Software coding the model:

QSARModel 4.0.4

Molcode Ltd., Turu 2, Tartu, 51014, Estonia

<http://www.molcode.com>

## 2. General information

### 2.1. Date of QMRF:

12.05.2010

### 2.2. QMRF author(s) and contact details:

Molcode model development team Molcode Ltd. Turu 2, Tartu, 51014, Estonia

[models@molcode.com](mailto:models@molcode.com) <http://www.molcode.com>

### 2.3. Date of QMRF update(s):

### 2.4. QMRF update(s):

### 2.5. Model developer(s) and contact details:

Molcode model development team Molcode Ltd. Turu 2, Tartu, 51014, Estonia

[models@molcode.com](mailto:models@molcode.com) <http://www.molcode.com>

### 2.6. Date of model development and/or publication:

12.05.2010

### 2.7. Reference(s) to main scientific papers and/or software package:

[1] Karelson M, Dobchev D, Tamm T, Tulp I, Jänes J, Tämm K, Lomaka A, Savchenko D & Karelson G (2008). Correlation of blood-brain penetration and human serum albumin binding with theoretical descriptors. ARKIVOC 16, 38-60.

[2] Karelson M, Karelson G, Tamm T, Tulp I, Jänes J, Tämm K, Lomaka A, Savchenko D and Dobchev D (2009), QSAR study of pharmacological permeabilities. ARKIVOC 2, 218-238.

### 2.8. Availability of information about the model:

Training and test sets are available.

### 2.9. Availability of another QMRF for exactly the same model:

None to date.

## 3. Defining the endpoint - OECD Principle 1

### 3.1. Species:

Not applicable

### 3.2. Endpoint:

1. Physical Chemical Properties 1.6. Octanol-water partition coefficient (Kow)

### 3.3. Comment on endpoint:

Annex (REACH) VII, test number 7.8 Partition coefficient

n-octanol/water, flask shake method

### **3.4.Endpoint units:**

Unitless

### **3.5.Dependent variable:**

logP

partition coefficient, the ratio of concentrations of a compound in two phases (water and octanol)

### **3.6.Experimental protocol:**

Partition coefficient was determined using the OECD Test Guideline 107 (shake flask method). The partition coefficient (P or Kow) is defined as the ratio of the equilibrium concentrations (C) of a dissolved substance in a two-phase system consisting of two largely immiscible solvents. In the case n-octanol and water:

$$P = Kow = C_{n-octanol}/C_{water}$$

The partition coefficient therefore is the quotient of two concentrations and is usually given in the form of its logarithm to base 10 (log P or log Kow).

The measuring range of the method is determined by the limit of detection of the analytical procedure. This should permit the assessment of values of log P in the range of -2 to 4 (occasionally when conditions apply, this range may be extended to log P up to 5) when the concentration of the solute in either phase is not more than 0,01 mol per litre [1-2].

The partitioning of organic compounds between aqueous and lipophilic phase is an important endpoint used extensively in medicinal chemistry, drug design, pharmacy, and environmental toxicity in predicting biological and hazardous effects of chemicals. No other physicochemical property has attracted as much interest in quantitative structure–activity relation (QSAR) studies as partition coefficient - lipophilicity (synonymously called hydrophobicity). The lipophilicity of the heterogeneous set of 223 compounds [3] is chosen from the compilation made by Hansch et al [ref 4; sect 9.2].

### **3.7.Endpoint data quality and variability:**

Experimental data from different sources were used. The previous modeling study [3] supports consistency of the data.

Statistics:

max value: 6.91

min value: -3.03

standard deviation: 1.64

skewness: -0.82

## **4.Defining the algorithm - OECD Principle 2**

### **4.1.Type of model:**

QSAR

### **4.2.Explicit algorithm:**

QSAR

Multilinear regression QSAR

$\text{LogP} = 2.48 - 2.61 \cdot \text{Average Bonding Information content (order 2)} + 2.13\text{E-}002 \cdot \text{Molecular surface area (AM1)} - 0.27 \cdot \text{count of H-acceptor sites (AM1)} - 0.25 \cdot \text{Charged (Zefirov) Surface Area of H atoms} - 0.12 \cdot \text{Charged (Zefirov) Surface Area of O atoms}$

#### **4.3.Descriptors in the model:**

- [1]Average Bonding Information content (order 2)
- [2]Molecular surface area (AM1) m2
- [3]count of H-acceptor sites (AM1)
- [4]Charged (Zefirov) Surface Area of H atoms m2
- [5]Charged (Zefirov) Surface Area of O atoms m2

#### **4.4.Descriptor selection:**

Initial pool of ~1000 descriptors. Stepwise descriptor selection based on a set of statistical selection rules:

- a) 1-parameter equations: Fisher criterion and R2 over threshold, variance and t-test value over threshold, intercorrelation with another descriptor not over threshold
- b) 2 parameter equations: intercorrelation coefficient below threshold, significant correlation with endpoint in terms of correlation coefficient and t-test).

Stepwise trial of additional descriptors not significantly correlated to any already in the model.

#### **4.5.Algorithm and descriptor generation:**

1D, 2D, and 3D theoretical calculations  
quantum chemical descriptors derived from AM1 calculation. Model developed by using multilinear regression.

#### **4.6.Software name and version for descriptor generation:**

QSARModel 4.0.4  
Molcode Ltd., Turu, 51014, Estonia  
<http://www.molcode.com>

#### **4.7.Chemicals/Descriptors ratio:**

39.2 (196 chemicals /5 descriptors)

### **5.Defining the applicability domain - OECD Principle 3**

#### **5.1.Description of the applicability domain of the model:**

By chemical identity: pesticides

By descriptor value range: this model is suitable for compounds that have descriptors in the following ranges:

Average Bonding Information content (order 2) (min: 0.28 , max: 1.22 ),  
Molecular surface area (AM1) (min: 65.02 , max: 525.10 ), count of H-acceptor sites (AM1) (min: 0, max: 9 ), Charged (Zefirov) Surface Area of H atoms (min: 0, max: 17.43 ), Charged (Zefirov) Surface Area of O atoms (min: 0 , max: 26.91).

#### **5.2.Method used to assess the applicability domain:**

Presence of functional groups in structures  
Range of descriptor values in training set with  $\pm 30\%$  confidence  
Descriptor values must fall between maximal and minimal descriptor

values of training set  $\pm 30\%$

### 5.3. Software name and version for applicability domain assessment:

QSARModel 4.0.4

Molcode Ltd., Turu 2, Tartu, 51014, Estonia

<http://www.molcode.com>

### 5.4. Limits of applicability:

## 6. Internal validation - OECD Principle 4

### 6.1. Availability of the training set:

Yes

### 6.2. Available information for the training set:

CAS RN: Yes

Chemical Name: Yes

Smiles: No

Formula: Yes

INChI: No

MOL file: Yes

### 6.3. Data for each descriptor variable for the training set:

All

### 6.4. Data for the dependent variable for the training set:

All

### 6.5. Other information about the training set:

196 data points: 14 negative; 182 positive values

### 6.6. Pre-processing of data before modelling:

### 6.7. Statistics for goodness-of-fit:

$R^2 = 0.76$  (Correlation coefficient);

$s = 0.82$  (Standard error of the estimate);

$F = 117.00$  (Fisher function);

### 6.8. Robustness - Statistics obtained by leave-one-out cross-validation:

$R^2_{cv} = 0.74$  LOO

### 6.9. Robustness - Statistics obtained by leave-many-out cross-validation:

$R^2_{cv} = 0.79$  LMO

### 6.10. Robustness - Statistics obtained by Y-scrambling:

### 6.11. Robustness - Statistics obtained by bootstrap:

### 6.12. Robustness - Statistics obtained by other methods:

ABC analysis (2:1 training : prediction) on sorted data divided into 3 subsets (A;B;C). Training set formed with 2/3 of the compounds (set A+B, A+C, B+C) and validation set consisted of 1/3 of the compounds (C, B, A)

average  $R^2(\text{fitting}) = 0.76$

average  $R^2(\text{prediction}) = 0.74$

## 7. External validation - OECD Principle 4

### 7.1. Availability of the external validation set:

Yes

**7.2.Available information for the external validation set:**

CAS RN: Yes

Chemical Name: Yes

Smiles: No

Formula: Yes

INChI: No

MOL file: Yes

**7.3.Data for each descriptor variable for the external validation set:**

All

**7.4.Data for the dependent variable for the external validation set:**

All

**7.5.Other information about the external validation set:**

21 data points: 1 negative values; 20 positive values

**7.6.Experimental design of test set:**

The full experimental dataset was sorted according to increasing values of logP and each tenth compound was assigned to the test set.

**7.7.Predictivity - Statistics obtained by external validation:**

$R^2=0.84$

**7.8.Predictivity - Assessment of the external validation set:**

The descriptors for the test set are in the limit of applicability.

**7.9.Comments on the external validation of the model:****8.Providing a mechanistic interpretation - OECD Principle 5****8.1.Mechanistic basis of the model:**

The solubility in octanol depends on the H bonding capability of the molecule, on the shape of the molecule and on the electrical charges that appear on the surface of the molecule. The descriptor count of H-acceptor sites (AM1) reflects the H bonding capability of the molecule and appears in the model with negative sign which means that the solubility in octanol decreases with increasing of the H bonding capability of the molecule. This is expected, because the H bonding capability renders the molecule more soluble in polar solvents like water than in non polar solvents like octanol. The presence of the descriptors Average Bonding Information Content (order 2) and Molecular Surface Area (AM1) in the model indicates that the solubility depends also on the shape and size of the molecule. The presence of charges on the surface of the molecule renders the molecule less soluble in octanol and more soluble in polar solvents like water. This can be seen from the presence in the model with negative sign of the descriptors Charged (Zefirov) Surface Area of H atoms and Charged (Zefirov) Surface Area of O atoms.

**8.2.A priori or a posteriori mechanistic interpretation:**

A posteriori mechanistic interpretation, consistent with published scientific interpretations of experiments [1,3-4]

**8.3.Other information about the mechanistic interpretation:**

Average Bonding Information Content (order 2) is a topological index that describes the atomic connectivity in the molecule and gives information about the shape and the size of the molecule.

The descriptors count of H-acceptor sites (AM1) reflects the H bonding capability of the molecule.

The descriptors Charged (Zefirov) Surface Area of H atoms and Charged (Zefirov) Surface Area of O atoms give information about the charges in the molecules localized at H and O atoms.

## 9. Miscellaneous information

### 9.1. Comments:

### 9.2. Bibliography:

- [1] Benfenati E, Gini G, Piclin N, Roncaglioni A & Vari MR (2003). Predicting logP of pesticides using different software. Chemosphere 53, 1155-1164. [http://dx.doi.org/10.1016/S0045-6535\(03\)00609-X](http://dx.doi.org/10.1016/S0045-6535(03)00609-X)
- [2] EU testing methods, Joint Research Centre <http://ecb.jrc.ec.europa.eu/testing-methods/>
- [3] Agrawal VK, Gupta M, Singh J & Khadikar PV (2005). A novel method of estimation of lipophilicity using distance-based topological indices: dominating role of equalized electronegativity. Bioorganic & Medicinal Chemistry 13, 2109–2120.
- [4] Hansch C, Leo A & Hockman D (1995). Exploring QSAR: Hydrolysis and Steric Constants. ACS Professional Reference Book. Washington, DC: The American Chemical Society.

### 9.3. Supporting information:

logP 2 training_196.sdf	<a href="http://qsardb.jrc.ec.europa.eu/qmrf/protocol/Q13-16-0061/attachment/A750">http://qsardb.jrc.ec.europa.eu/qmrf/protocol/Q13-16-0061/attachment/A750</a>
logP 2 test_21.sdf	<a href="http://qsardb.jrc.ec.europa.eu/qmrf/protocol/Q13-16-0061/attachment/A751">http://qsardb.jrc.ec.europa.eu/qmrf/protocol/Q13-16-0061/attachment/A751</a>

Test set(s)

## 10. Summary (JRC QSAR Model Database)

### 10.1. QMRF number:

Q13-16-0061

### 10.2. Publication date:

2013-07-02

### 10.3. Keywords:

Molcode;logP;octanol-water partition coefficient;pesticide;

### 10.4. Comments:

former Q19-10-30-300